

Cosmic flows in the nearby universe from Type Ia Supernovae

Stephen J. Turnbull¹, Michael J. Hudson^{1,2}, Hume A. Feldman³, Malcolm Hicken⁴, Robert P. Kirshner⁴, Richard Watkins⁵

¹ *Department of Physics and Astronomy, University of Waterloo, Waterloo, Ontario, N2L 3G1, Canada*

² *Perimeter Institute for Theoretical Physics, 31 Caroline St N, Waterloo, Ontario, N2L 2Y5, Canada*

³ *Department of Physics & Astronomy, University of Kansas, Lawrence, KS 66045, USA*

⁴ *Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138 USA*

⁵ *Department of Physics, Willamette University, Salem, OR 97301, USA*

E-mail: s4turnbu@uwaterloo.ca

8 November 2011

ABSTRACT

Peculiar velocities are one of the only probes of very large-scale mass density fluctuations in the nearby Universe. We present new “minimal variance” bulk flow measurements based upon the “First Amendment” compilation of 245 Type Ia supernovae (SNe) peculiar velocities and find a bulk flow of 249 ± 76 km s^{−1} in the direction $l = 319^\circ \pm 18^\circ$, $b = 7^\circ \pm 14^\circ$. The SNe bulk flow is consistent with the expectations of Λ CDM. However, it is also marginally consistent with the bulk flow of a larger compilation of non-SNe peculiar velocities (Watkins, Feldman, & Hudson 2009). By comparing the SNe peculiar velocities to predictions of the IRAS Point Source Catalog Redshift survey (PSCz) galaxy density field, we find $\Omega_m^{0.55} \sigma_{8,\text{lin}} = 0.40 \pm 0.07$, which is in agreement with Λ CDM. However, we also show that the PSCz density field fails to account for 150 ± 43 km s^{−1} of the SNe bulk motion.

1 INTRODUCTION

In the standard cosmological model, gravitational instability causes the growth of structure and peculiar velocities. In the regime where the perturbations are linear, there is a simple relationship between density and peculiar velocity (Peebles 1993):

$$\mathbf{v}(\mathbf{r}) = \frac{f}{4\pi} \int_0^\infty d^3\mathbf{r}' \delta(\mathbf{r}') \frac{\mathbf{r}' - \mathbf{r}}{|\mathbf{r}' - \mathbf{r}|^3} \quad (1)$$

where the growth factor f is equal to $\Omega_m^{0.55}$ in flat Λ CDM models (Linder 2005), δ is the normalized mass density fluctuation field, $\delta = (\rho - \bar{\rho})/\bar{\rho}$, and $\mathbf{r}$ are coordinates in units of km s^{−1}.

Given set of peculiar velocities, one can define a bulk flow as their average velocity; ideally the peculiar velocity tracers are dense and numerous enough that the resulting average is representative of the velocity of the volume. The bulk flow is then primarily due to structures on scales larger than the volume over which the bulk flow is measured (see Appendix A of Juszkiewicz et al. 1990 for a derivation). Hence, bulk flows are probes of the large-scale power spectrum of *matter* density fluctuations.

The Λ CDM model, once normalized by WMAP7 (Larson et al. 2011) observations of the Cosmic Microwave Background (CMB), fully specifies the r.m.s. fluctuations of δ on all scales, and hence the cosmic r.m.s. of bulk flows (Watkins et al. 2009). While most studies of bulk flows agree

on the general direction of the flow, there is some disagreement as to the amplitude and scale. Watkins et al. (2009) applied a “Minimal Variance” (MV) weighting scheme to a compilation of 4481 peculiar velocity measurements. Their results correspond to a sample with an effective Gaussian window of $50 h^{-1}$ Mpc and show a bulk flow of 407 ± 81 km s^{−1} towards $l = 287^\circ \pm 9^\circ$, $b = 8^\circ \pm 6^\circ$, which is in conflict with Λ CDM + WMAP7 at the 98 percent confidence level. The most controversial bulk flow result is the kinetic Sunyaev-Zeldovich flow dipole reported by Kashlinsky et al. (2010), who found a bulk flow on the order of 1000 km s^{−1} in the direction of $l = 296^\circ \pm 28^\circ$, $b = 39^\circ \pm 14^\circ$ over a scale of at least $800 h^{-1}$ Mpc. If correct, this result would strongly conflict with Λ CDM + WMAP7.

Another approach to understanding large-scale motions is to try to reconstruct the motion of the LG with respect to the CMB (627 ± 22 km s^{−1} towards $l = 276^\circ \pm 3^\circ$, $b = 30^\circ \pm 2^\circ$; Kogut et al. 1993) by measuring the distribution of galaxies and calculating the peculiar velocity of the Local Group (LG) using Eq. (1). Given the gravitational instability model of linear theory, the predicted velocity should converge to the measured CMB dipole for a sufficiently large survey volume. The application of Eq. (1) is difficult in practice because there are few redshift surveys that are both all-sky and deep. For example, Rowan-Robinson et al. (2000) found that the predicted dipole from the IRAS Point Source Catalog Redshift survey (Saunders et al. 2000, hereafter PSCz) converged to 13.4 degrees of the CMB dipole by

30,000 km s⁻¹. However, Basilakos & Plionis (2006) reanalyzed the same data set and found that significant power was required on large scales, which was missed by the original analysis. Other studies have been based on the Two Micron All-Sky Survey Redshift Survey (Skrutskie et al. 2006, hereafter 2MRS): Erdoğdu et al. (2006) and found probable convergence, but Lavaux et al. (2010) concluded that convergence was not obtained by 12,000 km s⁻¹, and may not be until well beyond 20,000 km s⁻¹. In another study, using only the infrared fluxes, Bilicki et al. (2011) concluded that even at an effective distance $\sim 300h^{-1}$ Mpc ($K_s < 13.5$) the flux dipole had not converged.

In this paper we use Type Ia SNe as our peculiar velocity tracers. SNe have also been used as peculiar velocity probes by a number of authors (Riess et al. 1995, 1997; Radburn-Smith et al. 2004; Pike & Hudson 2005; Lucey et al. 2005; Haugboelle et al. 2006; Colin et al. 2011; Dai et al. 2011).

An outline of this paper is as follows: In section 2, we introduce the data sets that were used. Section 3 presents the bulk flow of the SNe, using both simple weighting schemes as well as the “Minimal Variance” scheme of Watkins et al. (2009). Section 4 compares individual SNe peculiar velocities to the predictions of the IRAS PSCz density field. We discuss the implications of our results in Section 5, and present our conclusions in Section 6. Throughout, we adopt $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, and quote distances in units of km s⁻¹.

2 DATA AND CALIBRATION

In this study, three primary data sets of nearby SNe (with distances less than 20,000 km s⁻¹) are combined.

We refer to the first of these data sets as the ‘Old’ sample and it contains 106 SN the youngest being from 2002, drawn from two sources: Jha et al. (2007) and Hicken et al. (2009). Of the SNe in the ‘Old’ sample, 34 are from Jha et al. (2007). The remaining 72 SN in ‘Old’ are from Hicken et al. (2009). The second data set, which we refer to as ‘Hicken’, contains the remaining 113 SNe from Hicken et al. (2009) after cutting objects at distances larger than 20,000 km s⁻¹ and cutting two more objects (sn2007bz and sn2007ba) because they deviated by more than 3σ after the first round of fitting (as described below). The last set is the recently released data set from ‘The Carnegie Supernova Project’ (Folatelli et al. 2010, hereafter CSP), containing 28 SNe. Two of these objects were discarded due to our 20,000 km s⁻¹ distance cut, leaving 26 usable SNe. The CSP’s reported uncertainties only reflected the derived distance modulus residual spread. A second intrinsic uncertainty (σ_{SN}) in the magnitude of the SNe was added in quadrature by fitting a flow model and reducing the reduced chi-squared fit to 1.00. The intrinsic uncertainty was found to be 0.107 mag (slightly smaller than the 0.12 mag found by the CSP due to cuts and the additional free parameters of bulk flow). For further discussion of the light curve fitting, and consequences thereof, for the ‘Old’ and ‘Hicken’ data sets see Appendix A.

We combine these three sets to create a new sample that

we dub the ‘First Amendment’ (A1) compilation which we consider to be an extension to the ‘Constitution’ data set¹.

Where the observed SNe in the data sets were known to be contained within a cluster of galaxies, the redshift of the cluster was used for the observed velocity distance rather than the redshift of the supernova itself. Substituting cluster velocities for supernova velocities removes a significant source of thermal noise as objects in clusters can have a velocity rms of thousands of km s⁻¹. This process was applied to all three data sets. For galaxies not in clusters, the redshift of the host galaxy was used if the host galaxy redshift was recorded in NED, which occurred in all but two cases. For the remaining two cases we used the redshifts of the SNe. Galactic longitudes and latitudes for the Carnegie set were also taken from NED.

The A1 data set has a characteristic or uncertainty-weighted depth of 58 h^{-1} Mpc, where we define the characteristic depth to be:

$$r_* = \frac{\sum r/\sigma^2}{\sum 1/\sigma^2} \quad (2)$$

where σ is the total uncertainty in each SNe’s peculiar velocity and $\mathbf{r}$ is the coordinates in units of km s⁻¹.

In Fig. 1 we present our results, our raw data, and the bulk flow directions that other surveys have found in an Aitoff projection. In Fig. 2 we present the A1 data set in a Hubble Diagram divided into its three subsets. For all three data sets the intrinsic uncertainty of SNe’s is the dominant source of error. Thus, for all our SNe the percent error is approximately 6 percent of the measured distance, with the scatter for the ‘Old’ and ‘Hicken’ subsets being larger.

3 BULK FLOW

In this section we discuss the bulk flow, which is the simplest statistic that can be derived from a peculiar velocity survey.

3.1 Methods

We use two methods to measure the bulk flow. The first is a Maximum Likelihood (ML) method that minimizes the measurement uncertainties. The ML method is the traditional method used and we consider it in order to compare new results with previous ML results. However, ML methods have the disadvantage of returning the bulk flow of a specific sparse *sample* of peculiar velocity tracers rather than the bulk flow of a regular *volume*. Comparisons between ML results are complicated by the different spatial sampling. Instead, what is of greater interest is the bulk flow of a standardized volume. To estimate this, we calculate the “minimum variance” (MV) bulk flow as first introduced by Watkins et al. (2009).

¹ The ‘Old’ and ‘Hicken’ sets combined resemble very closely the ‘Constitution’ set from Hicken et al. (2009) in terms of which supernovae are included. The light curve fitter used here (MLCS2k2) differs from that of the ‘Constitution’ data set (SALT2).




Figure 1. An Aitoff projection of our data with circles (asterisks) representing SNe with peculiar velocities towards (away from) the LG. Larger symbols represent larger peculiar velocities in accordance with the scale shown top and bottom left. Also plotted in triangles are the direction motion of the LG with respect to the CMB, the Watkins et al. (2009) bulk flow direction, the kinetic Sunyaev-Zel’dovich bulk flow direction of Kashlinsky et al. (2010, labelled KAEK), and our new results (labeled SNBF for the bulk flow results from Section 3 and SNRF for the residual flow discussed in Section 4).




Figure 2. A Hubble diagram showing the three subsets that make up the A1 data set: ‘Old’ (red filled circles), ‘Hicken’ (Blue stars) and ‘Carnegie’ (green triangles). The error bars can be seen to be approximately constant in the log log diagram, or increasing proportionally to the distance, as to be expected with the dominant error in most cases being the intrinsic uncertainty in SNe.

3.1.1 Maximum Likelihood

In general, we fit a simple flow model (v_{pred}) to the SNIa peculiar velocity data. In the case of the bulk flow $\mathbf{V}$ in the CMB frame, this flow model reduces to the radial component of the bulk flow vector for each SNe, i. e.:

$$v_{pred,i} = \mathbf{V} \cdot \hat{\mathbf{r}}_i \quad (3)$$

where $\hat{\mathbf{r}}_i$ is the unit vector pointing to each supernova.

In the maximum likelihood method the weights are simply determined by the total uncertainty on the peculiar velocity of each object. Uncertainties in the observed peculiar velocity can be approximated well by a Gaussian, in which case the Maximum-Likelihood solution can be approximated by the following χ^2 :

$$\chi^2 = \sum_i \frac{[cz_{obs,i} - (r_i + v_{pred,i})]^2}{\sigma_i^2} \quad (4)$$

where cz_{obs} is the observed redshift in km s^{-1} , r_i is the distance converted from the reported distance modulus in km s^{-1} , $v_{pred,i}$ is the model velocity we are trying to measure as predicted for SNe i , and σ_i is the total uncertainty on the peculiar velocity of object i in units of km s^{-1} . This total uncertainty is the quadrature sum of the measurement error $\sigma_{m,i}^2$, the intrinsic uncertainty on SNe magnitude σ_{SN} (both converted from magnitudes to km s^{-1}) and a “thermal noise” term (σ_{th}) in units of km s^{-1} due to uncertainties in the flow model, such that:

$$\sigma_i^2 = \sigma_{m,i}^2 + \sigma_{SN}^2 + \sigma_{th}^2 \quad (5)$$

Note that since the σ_{SN}^2 term is converted from an uncertainty on magnitude, it is proportional to the distance to

the SN. σ_{SN}^2 is often the dominant source of uncertainty since the thermal term is only important in nearby supernovae where the $\sigma_{m,i}^2$ and σ_{SN}^2 terms are small. The results are only weakly dependant upon the precise value chosen. Here, where the flow model is a simple bulk flow, we set the thermal noise to 250 km s^{-1} , which is consistent with previous work. The impact of this choice for the thermal noise is discussed in the results below.

We let each sub-sample of the A1 data set have a freely-varying independent Hubble term to identify degeneracies, to avoid underestimating final uncertainties, and to account for the fact that each subset may have slightly different calibration. None of the fits preferred a Hubble value that varied by more than 1 percent from the value drawn from the original sources.

3.1.2 Minimum Variance Method

While the ML method described above is the best estimator of the bulk flow of a sparse sample, it is restricted in that it can only really characterize a particular survey, that will have its own errors and a specific and somewhat ill-defined geometry. The ML method is also, in a sense, density sampled, with higher density regions being more likely to contain a SN than voids. Most importantly, because the weights in the ML method are determined by the uncertainty on position in km s^{-1} , ML methods can be dominated by nearby SN that have smaller distance uncertainties.

To better approximate a *volume*-weighted bulk flow, we use the prescription described in Watkins et al. (2009) to estimate the volume flow. Each SN is weighted so as to minimize the variance between the bulk flow measured in the real sample and the bulk flow as it would be measured in a perfectly-sampled 3D Gaussian. We adopt a Gaussian with an “ideal” radius $R_I = 50h^{-1} \text{ Mpc}$. Effectively, weights are assigned to each SN based on their proximity to other SNe in the data set, and on how they compare with an ideal uniform sampling. This weighting scheme is specifically designed to maximize sensitivity to large scales. The MV weighting scheme has been tested using mock catalogues drawn from N-body simulations by Agarwal et al. 2011 (in preparation), who demonstrate that the recovered MV bulk flows are unbiased and have errors within the range expected from linear theory.

3.2 Consistency of SNe Subsamples

Before analysis of the combined SNe sample is undertaken, it is important to confirm that the data subsamples agree with one another. We calculate a χ^2 statistic for each pair of subsamples, following the analysis of Watkins et al. (2009), which accounts for sparse sampling effects. The χ^2 statistic we use is given by the equation,

$$\chi^2 = \sum_{i,j} (\Delta V_i)(C_{i,j})^{-1}(\Delta V_j). \quad (6)$$

where $\Delta \vec{V}$ is the bulk flow vector, and $\mathbf{C}$ is the covariance matrix taking into account the window functions of both surveys and the power spectrum, (see in equations 21 - 23 of Watkins et al. (2009)). The results are shown in Table 1. In summary, we find that all three subsamples are consistent with each other.

Table 1. χ^2 for 3 DoF for the surveys for $\Omega_m = 0.258$. If the χ^2 value is greater than 7.8, the two surveys disagree at a greater than 95% confidence level. The probabilities reported are the χ^2 test probability of agreement between the two.

Survey	$R_I = 50h^{-1} \text{ Mpc}$	
	χ^2	Probability
Old vs Hicken	0.173	98.2
Old vs Carnegie	2.293	51.4
Hicken vs Carnegie	1.369	71.3

3.3 Results

In Table 2, we present a summary of the results from the bulk flow, subdivided by data set and by weighting scheme. The ML bulk flow for the A1 sample was found to be $197 \pm 56 \text{ km s}^{-1}$ in direction $l = 295^\circ \pm 16^\circ$, $b = 11^\circ \pm 14^\circ$. This is significantly different from zero at the 99.9% confidence level.

As discussed above, the ML method gives most weight to SNe with the lowest errors in units of km s^{-1} , i.e. the nearest SNe. In order to reduce the impact of these nearby SNe, it is interesting to redetermine the bulk flow excluding nearby objects. The middle section of Table 2 shows the bulk flow using only SNe with $6000 \text{ km s}^{-1} < d < 20000 \text{ km s}^{-1}$. This subsample indicates a slightly higher amplitude flow, albeit with larger error bars: $330 \pm 120 \text{ km s}^{-1}$ towards $l = 321^\circ$, $b = 20^\circ$.

Finally, the MV results shown in the third section of Table 2 should give the most robust estimates of the flow of a Gaussian volume of radius $50 h^{-1} \text{ Mpc}$. For the entire A1 sample, the MV flow is $248 \pm 87 \text{ km s}^{-1}$ in the direction $l = 319^\circ \pm 25^\circ$, $b = 7^\circ \pm 13^\circ$.

These values are lower than the LG’s motion in the CMB frame, indicating that some of the LG’s motion must come from structures within our survey volume (such as the Virgo and Hydra-Centaurus superclusters).

To investigate the sensitivity of our results to the value of the thermal noise we adjusted it by $\pm 100 \text{ km s}^{-1}$; When so tested, the final magnitude of the A1 sample MV flow only changed by $\pm 31 \text{ km s}^{-1}$.

3.4 Bulk Flow: Cosmology and Comparisons

It is interesting to compare our ML bulk flow result to that of Colin et al. (2011), who apply a maximum likelihood bulk flow fit to the Union2 catalogue of Type Ia SNe (Amanullah et al. 2010). The Union2 catalogue contains 557 SNe, of which 165 are within $30,000 \text{ km s}^{-1}$. Colin et al. (2011)’s analysis yields a bulk flow velocity of $260 \pm 150 \text{ km s}^{-1}$ based on SNe within 18000 km s^{-1} . Our A1 sample yields a ML result of $196 \pm 55 \text{ km s}^{-1}$, which is consistent with theirs. It must be noted that the agreement between these results is not as significant as might at first be assumed because there is significant overlap between the data sets. However, Union2 uses SALT2 rather than MLCS2k2 ($R_V=1.7$) to obtain SN distances from the light-curve data.

Table 2. Bulk flow for all three SNIa data subsets and the combined First Amendment set. For comparison MV50 result from Watkins et al. (2009) are also included. Note the uncertainties quoted for the ML method are the just propagated uncertainties from measurements, the uncertainties for the MV method also include an approximation of additional noise due to non-uniform sampling.

	Number	Mag	l°	b°	V_X	V_Y	V_Z
ML Thermal noise=250 km s ⁻¹					km s ⁻¹	km s ⁻¹	km s ⁻¹
Old	106	226±76	307±21	4±15	136±76	-180±81	-35±59
Hicken	113	142±85	283±41	30±37	27±94	-120±96	71±78
Carnegie	26	260±140	330±170	76±38	54±182	-35±230	250±150
First Amendment	245	196±55	300±17	15±14	94±55	-165±58	50±44
ML Thermal noise=250 km s ⁻¹ , $d > 6,000$ km s ⁻¹							
Old	45	450±190	331±26	6±21	390±200	-210±190	44±160
Hicken	76	280±180	313±33	27±25	170±170	-180±190	130±110
Carnegie	15	1132±850	117±14	16±20	-490±540	970±810	310±300
First Amendment	136	330±120	321±20	16±15	250±120	-200±130	94±84
MV Weighting $R_I = 50h^{-1}$ Mpc Thermal noise=250 km s ⁻¹							
Old	113	240±110	318±26	-4±21	180±110	-160±110	-16±86
Hicken	113	250±110	310±25	5±20	160±110	-190±110	20±85
Carnegie	28	250±150	0±340	81±43	40±190	0±240	250±150
First Amendment	254	249±76	319±18	7±14	186±75	-162±77	32±59
Watkins et al. (2009)	4481	407±81	287±9	8±6	114±49	-387±53	57±37

Dai et al. (2011) also analyzed the Union2 catalog, splitting it into two subsets. They defined a nearby set with 132 SNe at $z < 0.05$ for which they found a bulk flow of 188^{+119}_{-103} km s⁻¹ towards $l = 290^{+39}_{-31}$, $b = 20^{+32}_{-32}$ which also agrees well with our results. The remaining 425 high- z SNe show no significant bulk flow. This is expected since the peculiar velocity errors are typically 6% of the distance to the source, and for this distant sample the errors per SNe measured in km s⁻¹ are extremely large.

Another interesting recent analysis of all peculiar velocities is by Watkins et al. (2009), who studied peculiar velocities mostly from Tully-Fisher, Fundamental Plane and SNe. They found that those subsamples had bulk flows consistent with each other². They combined the individual peculiar velocity samples into a “Composite” sample of 4481 peculiar velocity tracers, which was found to have a MV50 bulk flow of 407 ± 81 km s⁻¹ towards $l = 287^\circ \pm 9^\circ$, $b = 8^\circ \pm 6^\circ$. This result is inconsistent with Λ CDM at the 98% CL. However, their sample is not independent of ours. 103 of the 108 SNe which make up the “Old” subset of A1 are common to both A1 and Composite, although the latter takes SNe distances from Tonry et al. (2003). When all SNe data are removed from the ‘Composite’ data set, the two surveys become completely independent, and can be compared using the same formalism described in Section 3 of this paper and Section 5.1 of Watkins et al. (2009). We find that the ‘Composite excluding SNe’ MV50 bulk flow and the A1 MV50 results are consistent with each other, although the agreement is marginal: χ^2 of 6.4 for 3 directional degrees of freedom yields a 9 percent probability that the two results are consistent.

Lastly, these results can be compared directly to the

expectations for a Λ CDM universe. A plot showing the expectations of the one dimensional rms for perfectly sampled Gaussian sphere can be found in the top three plots of figure 5 from Feldman, Watkins, & Hudson (2010). The expectation for the one dimensional rms for a perfect Gaussian is 80 km s⁻¹. When you take into account the sparse sampling of the real A1 data set, this rises slightly to 91.2 km s⁻¹ assuming a σ_8 of 0.8 and Ω_m of 0.258.³ If you then include the propagation of measurement uncertainties the total expected rms for surveys equivalent to ours at different locations in space is 121 km s⁻¹. This prediction leads to χ^2 of 3.70 for 3 directional degrees of freedom yielding a 70 percent probability that the A1 data set is consistent with Λ CDM.

4 PREDICTED GRAVITY FIELD

4.1 Introduction

The MV weighting scheme discussed above is designed to suppress the effects of small-scale flows that would otherwise “alias” power into the bulk-flow statistic. An alternate method for removing the effects of small-scale structure on flow measurements is to assume gravitational instability and linear perturbation theory Eq. (1) and to predict the peculiar velocities using a model of the density field (derived from an all-sky galaxy redshift survey). The result is a model-dependent correction to measured peculiar velocities which can separate local effects from large-scale density waves from outside the survey volume.

³ In the rest of the paper because SN distances are insensitive to the value of Ω_m we used $\Omega_m = 0.3$. The predicted one dimensional rms drops slightly to 88.4 if this slightly higher value of Ω_m is used in the prediction

² Except for the BCG sample of Lauer & Postman (1994), which was excluded from further analysis.

Suppose we have an all-sky redshift survey that extends to a distance $R_{\max}$. We will model the peculiar velocity of a given SN located at position $\mathbf{r}$ by setting $v_{\text{pred},i}$ of Eq. (4) to a function with two terms:

$$\mathbf{v}_{\text{pred}}(\mathbf{r}) = \frac{\beta}{4\pi} \int_0^{R_{\max}} d^3\mathbf{r}' \delta_g(\mathbf{r}') \frac{\mathbf{r}' - \mathbf{r}}{|\mathbf{r}' - \mathbf{r}|^3} + \mathbf{U} \quad (7)$$

where $\beta = f/b$, b is the linear bias between galaxy density and mass density, $\mathbf{U}$ is the *residual* bulk flow of the volume being driven by mass structure beyond $R_{\max}$. In principle, the residual velocities have tidal (shear) and higher order terms. Feldman et al. (2010) measured the tidal and higher order terms for the “Composite” sample of Watkins et al. (2009), but found them to be small. We neglect these terms here and model the residual as a simple bulk flow $\mathbf{U}$.

The first term of v_{pred} is the predicted peculiar velocity induced by structure within the redshift survey volume ($r < R_{\max}$). The model is scaled by β to match the observed peculiar velocities of the SNe tracers. The peculiar velocity data therefore yields information about Ω_m and b . The residual bulk flow $\mathbf{U}$ is the additional velocity of the entire redshift survey volume in the CMB reference frame, and is presumably due to sources beyond $R_{\max}$. In an ideal survey, $\mathbf{U}$ would be completely independent of any structure within $R_{\max}$. This de-coupling of $\mathbf{U}$ from β means that $\mathbf{U}$ can be used to test consistency with $\Lambda\text{CDM} + \text{WMAP7}$ on large scales and β can do so on smaller scales.

4.2 Data and Method

The PSCz is both all-sky and deeper than, for example, the 2MRS (Skrutskie et al. 2006). Here we use the PSCz density field reconstructed by Branchini et al. (1999). For our study we applied the same $20,000 \text{ km s}^{-1}$ limit to the PSCz as we applied to the SNe. The PSCz density field in the Supergalactic Plane is shown in Fig. 3.

We fit the SNe data using the same method as in Section 3.1.1, but now with a new model as given by Eq. (7). Since the integral is specified by the PSCz density field, the free parameters are β and the three components of $\mathbf{U}$. Since the PSCz plus bulk flow is a better flow model than a simple bulk flow, we reduce the thermal component to 150 km s^{-1} , which is consistent with previous studies (Hudson et al. 2004).

4.3 Results

The results of the fits to each subset are given in Table 3. We find that the results from independent subsets are consistent with each other. For the A1 sample, the magnitude of the residual bulk flow was found to be $150 \pm 43 \text{ km s}^{-1}$ in direction $l = 345^\circ \pm 20^\circ$, $b = 8^\circ \pm 13^\circ$. This is significantly different from 0 at the 99.6% CL.

The value of β was found to be 0.53 ± 0.08 , and is shown in Fig. 4. The fit is sensitive to a single outlier, sn1992bh, for which the PSCz prediction is rather high (1719 km/s). Excluding this SN, we find $\beta = 0.57 \pm 0.08$.

Again we investigated the sensitivity of these results on the thermal noise term by changing it by $\pm 100 \text{ km s}^{-1}$. Again the magnitude off the flow only changed by $\pm 20 \text{ km s}^{-1}$, and the β changed by ± 0.03 .




Figure 3. The Supergalactic Plane. The PSCz galaxy density field is shown by the contours, predicted peculiar velocities as small black arrows, and measured supernova positions as “tadpoles” with dots showing measured positions and tails showing the magnitude of the measured radial peculiar velocity. The thick black contour corresponds to a $\delta = 0$ (or contours where the density is the mean universal density). The red(filled circles) SNe have peculiar velocities away from the LG and the blue(open circles) SNe have peculiar velocities towards the LG.




Figure 4. The observed peculiar velocity minus the measured bulk flow as a function of the linear-theory-predicted peculiar velocity for each SN, assuming $\beta = 1$. The circular symbol diameter scales with the inverse of the uncertainty (hence symbol area is proportional to weight). Representative error bars are shown in the top left. The slope is the fitted $\beta = 0.53$.

Table 3. Results from all three data subsets and the A1 full set with a linear perturbation theory model fit with a known matter distribution to fit β and the residual flow U . Fit with β as a free parameter and with a thermal noise of 150 km s^{-1} . Note the uncertainties quoted for this method are the just propagated uncertainties from the measurements.

	# of SN	Mag	l°	b°	U_X	U_Y	U_Z	β	χ^2	Dof
					km s^{-1}	km s^{-1}	km s^{-1}			
Old	106	190 ± 59	349 ± 22	0 ± 14	187 ± 60	-36 ± 73	0 ± 46	$.45 \pm .11$	139	101
Hicken	113	86 ± 77	347 ± 54	9 ± 41	84 ± 87	-19 ± 83	13 ± 66	$.62 \pm .13$	102	108
Carnegie	26	290 ± 150	347 ± 41	31 ± 26	240 ± 170	-50 ± 190	151 ± 130	$.82 \pm .33$	21	21
First Amendment	245	150 ± 43	345 ± 20	8 ± 13	144 ± 44	-38 ± 51	20 ± 35	$.53 \pm .08$	270	238

4.4 Gravity field: cosmology and comparisons

As noted above, the residual bulk flow $\mathbf{U}$ is significantly different from zero at the 99.6% confidence level. This means that there are structures not found in the PSCz catalogue that contribute significantly to the total peculiar velocity of the LG. As discussed in detail in Hudson et al. (2004) these structures could be missing from the PSCz because they are (i) outside the survey volume (ii) in the Zone Of Avoidance, or (iii) present but underrepresented. The latter scenario may arise because the IRAS (far-infrared) selection on which PSCz is based is sensitive to dusty spirals, but less so to the mostly dust-free early types. Lavaux & Hudson (2011), using the 6dF survey of 2MASS-selected galaxies, showed that the Shapley and Horologium-Reticulum superclusters generate significantly more peculiar velocity than predicted by the PSCz, even allowing for a different β for 2MASS galaxies.

We found that, for IRAS-selected galaxies, $\beta = 0.53 \pm 0.08$. This value of β_I is in good agreement with the IRAS average of 0.50 ± 0.02 reported by Pike & Hudson (2005), and with the SNIa-based result $\beta_I = 0.55 \pm 0.06$ of Radburn-Smith et al. (2004). Comparing fitted values of β between redshift surveys of different galaxy types is complicated by the fact that the bias factor b need not be the same because different galaxy types may trace the underlining mass density differently. This problem can be alleviated by noting that in linear theory the r.m.s. fluctuation of the survey galaxies, say in an $8 h^{-1} \text{ Mpc}$ top hat sphere ($\sigma_{8,\text{gal}}$) is proportional to the true matter r.m.s. fluctuations in a volume of the same size (i.e. $\sigma_{8,\text{gal}} = b\sigma_{8,\text{mass}}$). Thus with our measured β_I and the known $\sigma_{8,I}$ from the IRAS PSCz of 0.80 ± 0.05 (Hamilton & Tegmark 2002), we can calculate the degenerate parameter pair $f\sigma_8$ (where we dropped the subscript ‘mass’). Our value of β corresponds to $f\sigma_8 = 0.424 \pm 0.069$.

We can then compare our $f\sigma_8$ to other studies. Davis et al. (2011) compared the 2MRS density field and the SFI++ peculiar velocity data, and derived $f\sigma_8 = 0.31 \pm 0.05$. This is lower than our result, but not significantly so (1.5σ).

The $f\sigma_8$ parameter can also be derived from WMAP7 results. Recall that WMAP is observing fluctuations at an early epoch, when the perturbations were still well in the linear regime. To compare to WMAP7, we can convert our non-linear σ_8 into the equivalent linear value using the prescription of Juszkiewicz et al. (2010). If we assume an Ω_m of 0.272, our $\sigma_{8,\text{lin}}$ becomes 0.814 compared to its non-linear value of 0.867. Using this value of $\sigma_{8,\text{lin}}$, $f\sigma_{8,\text{lin}}$ drops to

0.40 ± 0.07 , which is in excellent agreement with the results of WMAP7 (Larson et al. 2011): $f\sigma_{8,\text{lin}} = 0.39 \pm 0.04$.

5 DISCUSSION

Attempts to determine the sources of the LG’s motion amount to determining the factors in Eq. (1). While early studies focussed on simple toy infall models, more recent studies have concentrated on models of the density field with the two free parameters β and $\mathbf{U}$. For a single object, such as the LG itself, there is a trade-off between these parameters. Lower values of β lead to larger values of $\mathbf{U}$, which are required in order to match the same $\mathbf{v}$ on the left-hand side of equation 7. This degeneracy can be broken with more than one measurement. We have shown that the PSCz does not account for all of the motion of the LG, although it is plausible that some of the missing signal comes from within the survey volume in the form of extra infall into the highest-density superclusters.

An alternative explanation for the bulk flow has been proposed, namely that the CMB temperature dipole, or part thereof, is intrinsic and does not represent the peculiar velocity of the LG (A “tilted” Universe: Turner 1991; Kashlinsky et al. 2008; Ma et al. 2011). This would lead to an illusory “bulk flow” which would extend well beyond the local volume, indeed to the horizon. The apparent $1005 \pm 267 \text{ km s}^{-1}$ bulk flow of $z < 0.25$ clusters claimed by Kashlinsky et al. (2010), which is well outside the expectations of ΛCDM bulk flows, might be explained by such an effect. In such a scenario, there is an additional “bulk flow” $\mathbf{U}_{\text{tilt}}$ which never vanishes no matter how deep a redshift survey R_{max} is used in Eq. (7). Our measured $\mathbf{U}$ thus provides an upper limit on the $\mathbf{U}_{\text{tilt}}$. The amplitude of the bulk flow found by Kashlinsky et al. (2010) is inconsistent with our measurement of $\mathbf{U} = 150 \pm 43 \text{ km s}^{-1}$. However, amplitude of the Kashlinsky et al. (2010) bulk flow is systematically uncertain. If we compare only the direction of the A1 fit $l = 345^\circ \pm 20^\circ$, $b = 8^\circ \pm 13^\circ$ and the Kashlinsky et al. (2010) direction $l = 296^\circ \pm 29^\circ$, $b = 39^\circ \pm 15^\circ$, the results are marginal: they disagree at approximately the 90% CL. Thus our results do not support the high amplitude bulk flow found by Kashlinsky et al. (2010).

6 CONCLUSION

We have analyzed the peculiar velocities of a 245 SNe dataset dubbed the “First Amendment”. Overall, we have

found that this new compilation is in marginal agreement with previous bulk flow results and is not in significant conflict with Λ CDM + WMAP7 predictions. The First Amendment compilation yields a bulk flow of $248 \pm 87 \text{ km s}^{-1}$ in the direction $l = 319^\circ \pm 25^\circ$, $b = 7^\circ \pm 13^\circ$.

We have compared the peculiar velocities to the predictions from the IRAS PSCz and have found $\Omega_m^{0.55} \sigma_{8,\text{lin}}$ of 0.40 ± 0.07 , which is in excellent agreement with the Λ CDM + WMAP7 predictions and other previous measurements.

A residual flow of $150 \pm 43 \text{ km s}^{-1}$ $l = 345^\circ \pm 20^\circ$, $b = 8^\circ \pm 13^\circ$ was found for the IRAS Point Source Catalog as normalized with the First Amendment SNe. This may suggest that the IRAS PSCz undersamples massive dense superclusters such as the Shapley Concentration. Nevertheless, the small amplitude of the residual flow is in conflict with “tilted Universe” scenarios such as might be favoured by the kSZ analysis of Kashlinsky et al. (2010).

As its name suggests, the First Amendment compilation is readily extendible as new SNe are found and their distances are published. Ongoing surveys such as CfA4 (95 SNe, Hicken, private communication), LOSS (Ganeshalingam et al. 2011), Palomar Transit Factory (Law et al. 2009), and CSP (Stritzinger et al. 2011, 50 more distances expected soon), and upcoming surveys such as SkyMapper (100 SNe per year with $z < 0.085$, Keller et al. 2007), Pan-Starrs, and LSST will eventually provide sufficient SNe to reduce the $20,000 \text{ km s}^{-1}$ bulk and residual flow uncertainties to the systematic limits. Future results on $f\sigma_{8,\text{lin}}$ are expected based on predicted peculiar velocities from the 2M++ redshift compilation (Lavaux & Hudson 2011). Additionally, although individually less precise, Fundamental Plane distances and peculiar velocities can contribute significant precision to bulk flow surveys by sheer numbers. We wish to re-analyze the full ‘Composite’ data set from Feldman & Watkins (2008) after replacing the 103 SNe currently contained in that data set with the 245 SNe of A1, as well as to add the Fundamental Plane peculiar velocities from NFPS (Smith et al. 2006) and 6dF (Jones et al. 2004) when they become available. For now the results are data-limited, but the future promises many fruitful results from many promising surveys, and we await them eagerly.

7 ACKNOWLEDGEMENTS

We thank Alex Conley for helpful comparisons of SNe data, and acknowledge useful discussions with John Lucey, Roya Mohayaee and Jacques Colin. SJT and MJH acknowledge the financial support of NSERC. RPK acknowledges support from the NSF grant AST-0907903.

This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Amanullah R. et al., 2010, *ApJ*, 716, 712
 Basilakos S., Plionis M., 2006, *MNRAS*, 373, 1112
 Bilicki M., Chodorowski M., Mamon G. A., Jarrett T., 2011, *ArXiv e-prints*
 Branchini E. et al., 1999, *MNRAS*, 308, 1
 Colin J., Mohayaee R., Sarkar S., Shafieloo A., 2011, *MNRAS*, 414, 264
 Dai D.-C., Kinney W. H., Stojkovic D., 2011, *JCAP*, 4, 15
 Davis M., Nusser A., Masters K. L., Springob C., Huchra J. P., Lemson G., 2011, *MNRAS*, 413, 2906
 Erdoğan P. et al., 2006, *MNRAS*, 368, 1515
 Feldman H. A., Watkins R., 2008, *MNRAS*, 387, 825
 Feldman H. A., Watkins R., Hudson M. J., 2010, *MNRAS*, 407, 2328
 Folatelli G. et al., 2010, *AJ*, 139, 120
 Ganeshalingam M., Li W., Filippenko A. V., 2011, *ArXiv e-prints*
 Hamilton A. J. S., Tegmark M., 2002, *MNRAS*, 330, 506
 Haugboelle T., Hannestad S., Thomsen B., Fynbo J., Sollerman J., Jha S., 2006, *ArXiv Astrophysics e-prints*
 Hicken M., Wood-Vasey W. M., Blondin S., Challis P., Jha S., Kelly P. L., Rest A., Kirshner R. P., 2009, *The Astrophysical Journal*, 700, 1097
 Hudson M. J., Smith R. J., Lucey J. R., Branchini E., 2004, *MNRAS*, 352, 61
 Jha S., Riess A. G., Kirshner R. P., 2007, *ApJ*, 659, 122
 Jones D. H. et al., 2004, *MNRAS*, 355, 747
 Juszkievicz R., Feldman H. A., Fry J. N., Jaffe A. H., 2010, *JCAP*, 2, 21
 Juszkievicz R., Vittorio N., Wyse R. F. G., 1990, *ApJ*, 349, 408
 Kashlinsky A., Atrio-Barandela F., Ebeling H., Edge A., Kocevski D., 2010, *ApJL*, 712, L81
 Kashlinsky A., Atrio-Barandela F., Kocevski D., Ebeling H., 2008, *ApJL*, 686, L49
 Keller S. C. et al., 2007, *Publications of the Astronomical Society of Australia*, 24, 1
 Kogut A. et al., 1993, *ApJ*, 419, 1
 Larson D. et al., 2011, *ApJ Supp*, 192, 16
 Lauer T. R., Postman M., 1994, *ApJ*, 425, 418
 Lavaux G., Hudson M. J., 2011, *ArXiv e-prints*
 Lavaux G., Tully R. B., Mohayaee R., Colombi S., 2010, *ApJ*, 709, 483
 Law N. M. et al., 2009, *PASP*, 121, 1395
 Linder E. V., 2005, *PRD*, 72, 043529
 Lucey J., Radburn-Smith D., Hudson M., 2005, in *ASP Conf. Ser. 329: Nearby Large-Scale Structures and the Zone of Avoidance*, pp. 21–26
 Ma Y.-Z., Gordon C., Feldman H. A., 2011, *Phys Rev D*, 83, 103002
 Peebles P. J. E., 1993, *Principles of Physical Cosmology*. Princeton University Press
 Pike R. W., Hudson M. J., 2005, *ApJ*, 635, 11
 Radburn-Smith D. J., Lucey J. R., Hudson M. J., 2004, *MNRAS*, 355, 1378
 Riess A. G., Davis M., Baker J., Kirshner R. P., 1997, *ApJL*, 488, L1+
 Riess A. G., Press W. H., Kirshner R. P., 1995, *ApJL*, 445, L91
 Rowan-Robinson M. et al., 2000, *MNRAS*, 314, 375
 Saunders W. et al., 2000, *MNRAS*, 317, 55
 Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
 Smith R. J., Hudson M. J., Lucey J. R., Nelan J. E., Wegner G. A., 2006, *MNRAS*, 369, 1419

- Stritzinger M. et al., 2011, ArXiv e-prints
 Tonry J. L. et al., 2003, ApJ, 594, 1
 Turner M. S., 1991, Phys Rev D, 44, 3737
 Watkins R., Feldman H. A., Hudson M. J., 2009, MNRAS, 392, 743

APPENDIX A: LIGHT CURVE PARAMETER COMPARISONS

The A1 data set is composed of three different SNe catalogues; this complicates the description of the light curve fitting procedures used because the catalogues used different methods. In the ‘Old’ sample are 34 SNe from Jha et al. (2007); most of which are fit using the MLCS2k2 light curve fitter with a reddening law parameter R_V of 3.1 (for SN with high extinction, R_V was a free fit parameter with a tight prior of 3.1). The remaining 72 SN in ‘Old’ are from Hicken et al. (2009) and are also fit using MLCS2k2, but with a reddening law parameter R_V of 1.7. The second data set, which we refer to as ‘Hicken’, contains the remaining 113 SNe from Hicken et al. (2009) and are all fit with MLCS2k2 with a reddening law parameter R_V of 1.7. The ‘Carnegie’ set containing 28 SNe were fit with a R_V as a free variable. The light curve fitter used for the ‘Carnegie’ set is described in detail in the original paper Saunders et al. (2000).

For SNe fit by Hicken et al. (2009), four distances were reported for each SN. We use the distances reported using the MLCS2k2 fitting procedure rather than either of the SALT procedures for multiple reason. To start the MLCS2k2 process determines host reddening on a case-by-case basis. Furthermore, of the two published MLCS2k2 methods we use the results with a reddening law parameter R_V of 1.7 instead of 3.1 since Hicken et al. (2009) show that the Hubble residuals for high-extinction SNIa’s using $R_V=3.1$ are systematically negative, (suggesting that the extinction is overestimated). We study the effect the choice of R_V parameter has upon bulk flow measurements to explore systematics. Hicken et al. (2009) provide distances to 162 SNe using both $R_V = 1.7$ and $R_V = 3.1$. We fit both of these data sets for bulk flows using the ML method to investigate the systematics. The results of this comparison can be seen in table A1. Although the results for the two light curve fitters agree to less than one σ in each of the 3 degrees of freedom, the data sets are fit to the same light curves, so they are not independent. This result highlights how large the systematic errors are for bulk flow surveys, in part reflected by the large σ_{SN} , which in most cases dominates the uncertainty budget for peculiar velocity surveys.

Table A1. Results for 162 SNe from Hicken et al. (2009) fit with the MLCS2k2 light curve fitter either with $R_V = 1.7$ or $R_V = 3.1$.

	Number	Mag	l°	b°	V_X	V_Y	V_Z
					km s ⁻¹	km s ⁻¹	km s ⁻¹
ML Thermal noise=250 km s ⁻¹							
$R_V = 1.7$	162	220±70	298±18	9±14	103±68	-191±73	35±52
$R_V = 3.1$	162	175±70	310±25	14±18	108±70	-131±75	43±53